

Meiosis

Formation of Gametes
(Eggs & Sperm)

Facts About Meiosis

- ✓ Preceded by **interphase** which includes **chromosome replication**
- ✓ Two meiotic divisions --- **Meiosis I and Meiosis II**
- ✓ Called **Reduction-division**
- ✓ Original cell is **diploid (2n)**
- ✓ Four daughter cells produced that are **monoploid (1n)**

Facts About Meiosis


- ✓ Daughter cells contain **half the number of chromosomes** as the original cell
- ✓ **Produces gametes** (eggs & sperm)
- ✓ **Occurs in the testes** in males (Spermatogenesis)
- ✓ **Occurs in the ovaries** in females (Oogenesis)

More Meiosis Facts

- ✓ Start with 46 double stranded chromosomes ($2n$)
- ✓ After 1 division - 23 double stranded chromosomes (n)
- ✓ After 2nd division - 23 single stranded chromosomes (n)
- ✓ Occurs in our germ cells that produce gametes

Why Do we Need Meiosis?

- ✓ It is the fundamental basis of sexual reproduction
- ✓ Two haploid ($1n$) gametes are brought together through fertilization to form a diploid ($2n$) zygote






What happens during meiosis I?

- ▣ As previously mentioned, the first round of nuclear division that occurs during the formation of gametes is called **meiosis I**. It is also known as the **reduction division** because it results in cells that have half the number of chromosomes as the parent cell.
- ▣ **Meiosis I** consists of four phases: **prophase I, metaphase I, anaphase I, and telophase I**.

Replication of Chromosomes

- ✓ Replication is the process of duplicating a chromosome
- ✓ Occurs prior to division
- ✓ Replicated copies are called sister chromatids
- ✓ Held together at centromere

Occurs in Interphase



Prophase I

- ▣ During prophase I, the chromosomes condense and become visible inside the nucleus. *Because each chromosome was duplicated during the S phase that occurred just before prophase I,* each now consists of two sister chromatids joined at the centromere. This arrangement means that each chromosome has the shape of an X.


- ▣ Once this chromosomal condensation has occurred, the members of each chromosome pair *(called homologous chromosomes, because they are similar in size and contain similar genes)*, align next to each other.
- ▣ At this point, the two chromosomes in each pair **become tightly associated** with each other along their lengths in a process called **synapsis**.

- ▣ Then, while the **homologous chromosomes** are tightly paired, the members of each pair **trade** adjacent bits of DNA in a process called **crossing over**, also known **as recombination**. This trading of genetic material creates **unique chromosomes that contain new combinations of alleles**.

Tetrads Form in Prophase I

Homologous chromosomes
(each with sister chromatids)


Join to form a TETRAD




Called Synapsis

Crossing-Over

- ✓ Homologous chromosomes in a tetrad cross over each other
- ✓ Pieces of chromosomes or genes are exchanged
- ✓ Produces Genetic recombination in the offspring



Crossing-Over




Crossing-over multiplies the already huge number of different gamete types produced by independent assortment

- ▣ At the end of prophase I, the nuclear membrane finally begins to break down. Outside the nucleus, the spindle grows out from centrosomes on each side of the cell.
- ▣ As in mitosis, the microtubules of the spindle are responsible for moving and arranging the chromosomes during division.

Metaphase I


- ▣ At the start of metaphase I, microtubules emerge from the spindle and attach to the kinetochore near the centromere of each chromosome.

- ▣ In particular, microtubules from one side of the spindle attach to one of the chromosomes in each homologous pair, while microtubules from the other side of the spindle attach to the other member of each pair.
- ▣ With the aid of these microtubules, the chromosome pairs *then line up along the equator of the cell, termed the metaphase plate (Figure 2).*



- ▣ Figure 2: Near the end of metaphase I, the homologous chromosomes align on the metaphase plate

Metaphase I




Homologous pairs of chromosomes align along the equator of the cell

Anaphase I


- During **anaphase I**, the microtubules disassemble and contract; this, in turn, separates the homologous chromosomes such that the two chromosomes in each pair are pulled toward opposite ends of the cell .
- This separation means that each of the daughter cells that results from meiosis I will have half the number of chromosomes of the original parent cell after interphase.
- Also, the sister chromatids in each chromosome still remain connected. As a result, each chromosome maintains its X-shaped structure.





- ▣ Figure 3: During anaphase I, the homologous chromosomes are pulled toward opposite poles of the cell.

Anaphase I




Homologs separate and move to opposite poles.

Sister chromatids remain attached at their centromeres.


Telophase I

- ▣ As the new chromosomes reach the spindle during **telophase I**, the cytoplasm organizes itself and divides in two.
- ▣ There are now two cells, and each cell contains half the number of chromosomes as the parent cell.
- ▣ In addition, the two daughter cells are not genetically identical to each other because of the recombination that occurred during prophase I .



- ▣ Figure 4: Telophase I results in the production of two nonidentical daughter cells, each of which has half the number of chromosomes of the original parent cell.

Telophase I




Nuclear envelopes reassemble.

Spindle disappears.

Cytokinesis divides cell into two.

Meiosis I: Reduction Division



Early Prophase I
(Chromosome
number doubled)

Late
Prophase I

Metaphase I

Anaphase I

Telophase I
(diploid)

Interkinesis

- ▣ At this point, the first division of meiosis is complete. The cell now rests for a bit before beginning the second meiotic division. During this period, **called interkinesis**, the nuclear membrane in each of the two cells reforms around the chromosomes. In some cells, the spindle also disintegrates and the chromosomes relax (although most often, the spindle remains intact).
- ▣ It is important to note, however, that no chromosomal duplication occurs during this stage.




What happens during meiosis II?

- ▣ *During meiosis II, the two cells once again cycle through four phases of division.*
- ▣ Meiosis II is sometimes referred to as an **equational division** because it does not reduce chromosome number in the daughter cells — rather, the daughter cells that result from meiosis II have the same number of chromosomes as the "parent" cells that enter meiosis II.


Prophase II

- ▣ As **prophase II** begins, the chromosomes once again condense into tight structures, and the nuclear membrane disintegrates. In addition, if the spindle was disassembled during interkinesis, it reforms at this point in time.

Prophase II



Nuclear envelope fragments.




Spindle forms.

Metaphase II


- ▣ The events of **metaphase II** are similar to those of mitotic metaphase — in both processes, the chromosomes line up along the cell's equatorial plate, also called the metaphase plate, in preparation for their eventual separation (Figure 5).



Metaphase II




Chromosomes align along **equator** of cell.




Anaphase II

- ▣ During **anaphase II**, microtubules from each spindle attach to each sister chromatid at the kinetochore. The sister chromatids then separate, and the microtubules pull them to opposite poles of the cell. As in mitosis, each chromatid is now considered a separate chromosome (Figure 6).
- ▣ This means that the cells that result from meiosis II will have the same number of chromosomes as the "parent" cells that entered meiosis II.




- ▣ Figure 6: Anaphase II involves separation of the sister chromatids

Anaphase II




Sister chromatids separate and move to opposite poles.




Telophase II

- ▣ Finally, in **telophase II**, nuclear membranes reform around the newly separated chromosomes, which relax and fade from view. As soon as the cytoplasm divides, meiosis is complete. There are now four daughter cells — two from each of the two cells that entered meiosis II — and each daughter cell has half the normal number of chromosomes (Figure 7). Each also contains new mixtures of genes within its chromosomes, thanks to recombination during meiosis I.




- ▣ Figure 7: Telophase II results in the production of four daughter cells

Telophase II



Meiosis II: Reducing Chromosome Number



Why is meiosis important?

- ▣ Meiosis is important because it ensures that all organisms produced via sexual reproduction contain the correct number of chromosomes. Meiosis also produces genetic variation by way of the process of recombination. Later, this variation is increased even further when two gametes unite during fertilization, thereby creating offspring with unique combinations of DNA. This constant mixing of parental DNA in sexual reproduction helps fuel the incredible diversity of life on Earth.



- ▣ Like mitosis, **meiosis** is a form of eukaryotic cell division. However, these two processes distribute genetic material among the resulting daughter cells in very different ways.
- ▣ **Mitosis** creates two identical daughter cells that each contain the same number of chromosomes as their parent cell.

- ▣ In contrast, **meiosis** gives rise to four unique daughter cells, each of which has half the number of chromosomes as the parent cell.
- ▣ Because **meiosis** creates cells that are destined to become **gametes** (or reproductive cells), this reduction in chromosome number is critical — without it, the union of two gametes during fertilization would result in offspring with twice the normal number of chromosomes!

- ▣ Apart from this reduction in chromosome number, **meiosis differs from mitosis** in yet another way.
- ▣ Specifically, meiosis creates new combinations of genetic material in each of the four daughter cells. These new combinations result from the exchange of DNA between paired chromosomes. Such exchange means that the gametes produced through meiosis exhibit an amazing range of genetic variation.

- ▣ Finally, unlike mitosis, meiosis involves two rounds of nuclear division, not just one. Despite this fact, many of the other events of meiosis are similar to those that occur in mitosis. For example, prior to undergoing meiosis, a cell goes through an interphase period in which it grows, replicates its chromosomes, and checks all of its systems to ensure that it is ready to divide.

- ▣ Like **mitosis**, meiosis also has distinct stages called prophase, metaphase, anaphase, and telophase. A key difference, however, is that during meiosis, each of these phases occurs twice — once during the first round of division, called meiosis I, and again during the second round of division, called meiosis II

Comparison of Divisions

	Mitosis	Meiosis
Number of divisions	1	2
Number of daughter cells	2	4
Genetically identical?	Yes	No
Chromosome #	Same as parent	Half of parent
Where	Somatic cells	Germ cells
When	Throughout life	At sexual maturity
Role	Growth and repair	Sexual reproduction

Questions